

# UCII Supplementation

## Humans Studies

1. Bagchi et al. Effects of orally administered undenatured type II collagen against arthritic inflammatory diseases: A Mechanistic exploration. *Int J Clin Pharmacol Res.* 2002; 22(34): 101–10.
2. Crowley et al. Safety and efficacy of undenatured type II collagen in the treatment of osteoarthritis of the knee; a clinical trial. *Int J Med Sci.* 2009; 6(6): 312–321.
3. Lugo et al. Undenatured type II collagen (UC-II<sup>®</sup>) for joint support: a randomized, double-blind, placebo-controlled study in healthy volunteers. *J Int Soc Sports Nutr.* 2013; 10: 48
4. Lugo et al. Efficacy and tolerability of an undenatured type II collagen supplement in modulating knee osteoarthritis symptoms: a multicenter randomized, double-blind, placebo-controlled study. *Nutr J.* 2016; 15: 14.
5. Mehra et al. A non-interventional, prospective, multicentric real life Indian study to assess safety and effectiveness of un-denatured type 2 collagen in management of osteoarthritis. *Int J Res Orthop.* 2019; 5(2): 315–320
6. Mohammed Abdul Azeem et al. The Study of undenatured type II collagen in the knee osteoarthritis *International Journal of Orthopaedics Traumatology & Surgical Sciences*, 2019; 5(01): 172–175
7. Yatish et al. Evaluation of clinical efficacy of undenatured type ii collagen in the treatment of osteoarthritis of knee. A randomized controlled study. *Int J Orthopaedics Sci.* 2020; 6(2): 497–500
8. Marone et al. Safety and toxicological evaluation of undenatured type II collagen. *Toxicol MechMethods.* 2010; 1–15

## Animal Studies

1. D'Altilio et al. Therapeutic Efficacy and Safety of Undenatured Type II Collagen Singly or in Combination with Glucosamine and Chondroitin in Arthritic Dogs. *Toxicol MechMethods.* 2007; 17(4): 189–96.
2. Peal et al. Therapeutic efficacy and safety of undenatured type-II collagen (UC-II) alone or in combination with (-)-hydroxycitric acid and chromemate in arthritic dogs. *J Vet Pharmacol Ther.* 2007 Jun; 30(3): 275–8
3. Deparle et al. Efficacy and safety of glycosylated undenatured type-II collagen (UC-II) in therapy of arthritic dogs. *J Vet Pharmacol Ther.* 2005; 28: 385–390
4. Bagchi et al. Efficacy and safety of undenatured type II collagen (UC-II) in arthritic horses. *Toxicol Lett.* 2007; 172: S223
5. Gupta et al. Comparative therapeutic efficacy and safety of type-II collagen (UC-II), glucosamine and chondroitin in arthritic dogs: pain evaluation by ground force plate. *J Anim Physiol Anim Nutr (Berl).* 2012; 96(5): 770–777.
6. Bagchi et al. Suppression of arthritic pain in dogs by undenatured type-II collagen (UC-II) treatment quantitatively assessed by ground force plate *Toxicol Lett.* 2009, 189, S231.
7. Gupta et al. Comparative therapeutic efficacy and safety of type-II collagen (UC-II), glucosamine and chondroitin in arthritic dogs: pain evaluation by ground force plate. *J Anim Physiol Anim Nutr (Berl).* 2012; 96(5): 770–777.
8. Bagi et al. Oral administration of undenatured native chicken type II collagen (UC-II) diminished deterioration of articular cartilage in a rat model of osteoarthritis (OA). *Osteoarthritis Cartilage.* 2017; 25(12): 2080–2090.
9. Blair et al. Palatability and tolerability of a novel joint supplement in the cat. *J Feline Med Surg.* 2017; 19: 961–969
10. Martello et al. Evaluation of The Efficacy of a Dietary Supplement in Alleviating Symptoms in Dogs with Osteoarthritis. *J Food Nutr Res.* 2017; 4(104): 1–8
11. Stabile et al. Evaluation of the Effects of Undenatured Type II Collagen (UC-II) as Compared to Robenacoxib on the Mobility Impairment Induced by Osteoarthritis in Dogs. *Vet Sci.* 2019, 6, 72.